

Revisiting the dynamical exponent equality $z = d$ for the dirty boson problem

Peter B. Weichman¹ and Ranjan Mukhopadhyay²

¹*BAE Systems, Advanced Information Technologies,
6 New England Executive Park, Burlington, MA 01803*

²*Department of Physics, Clark University, Worcester, MA 01610*

It is shown that previous arguments leading to the equality $z = d$ (d being the spatial dimensionality) for the dynamical exponent describing the Bose glass to superfluid transition may break down, as apparently seen in recent simulations (Ref. [8]). The key observation is that the major contribution to the compressibility, which remains finite through the transition and was predicted to scale as $\kappa \sim |\delta|^{(d-z)\nu}$ (where δ is the deviation from criticality and ν is the correlation length exponent) comes from the analytic part, not the singular part of the free energy, and therefore is not restricted by any conventional scaling hypothesis.

PACS numbers: 64.60.Fr, 67.40.-w, 72.15.Rn, 74.78.-w,

Beginning with the realization that the zero temperature onset transition of superfluidity in a random medium should properly be treated as a fluctuation driven quantum phase transition [1, 2, 3, 4], a number of scaling arguments have been put forward [2, 3] to determine or to place bounds on the critical exponents describing it. Probably the most significant of these was an argument that the dynamical exponent, z , describing the relative divergence of the temporal and spatial correlation lengths via $\xi_\tau \sim \xi^z$, should be equal to the dimension, d , of space. The argument was supported by exact calculations in $d = 1$, where indeed $z = 1$ [3], by $1 + \epsilon$ renormalization group calculations [5] which, however, are not completely rigorous due to the absence of a form for the Hamiltonian for noninteger $d > 1$, and by a series of quantum Monte Carlo (QMC) studies in $d \leq 2$ [6, 7, 8]. Results in $d > 2$ are restricted to a badly controlled double ϵ -expansion which is inappropriate for testing an exact scaling argument of this type [4, 9], but nevertheless points to values of z significantly larger than unity [9].

With one controversial exception [7], the earlier QMC results [6] were fit to $z = d$, but the numerical error bars were not very tight. The more recent QMC study in $d = 2$ [8] provided a much more stringent test: using larger systems and a joint, optimal critical functional form fit to three separate thermodynamic quantities, the value $z = 1.40 \pm 0.02$ was found, strongly violating $z = d$. In light of this, the purpose of this paper is to revisit the scaling arguments [2, 3] that led to $z = d$, and show that they can in fact *break down*: z might be independent of all the other exponents. In the absence of further theoretical justification, we raise the possibility that z is unconstrained by any simple scaling argument.

Before presenting details, we briefly review the essence of the scaling arguments and indicate where they could go wrong. The helicity modulus, Υ [or superfluid density, $\rho_s = (m^2/\hbar^2)\Upsilon$], quantifies the response of the superfluid to gradients in the phase, ϕ , of the order parameter. Specifically, the free energy density contains a correction $\Delta f_x = (\Upsilon/2\beta V) \int_0^\beta d\tau \int d^d x |\nabla \phi(\mathbf{x}, \tau)|^2$, where V

is the volume, $\beta = 1/k_B T$, and τ is the usual imaginary time variable. The essence of the Josephson scaling argument is that since $\nabla \phi$ has dimensions of inverse length, it should scale as $\xi^{-1} \sim |\delta|^\nu$ where δ is the deviation of the control parameter from criticality and ν is the correlation length exponent. On the other hand the singular part of the free energy, f_s , is defined to scale as $|\delta|^{2-\alpha}$. This implies that $\Upsilon \sim |\delta|^\nu$ with $\nu = 2 - \alpha - 2\nu = (d + z - 2)\nu$, where the last equality follows from the quantum hyperscaling relation $2 - \alpha = (d + z)\nu$ [3]. Now, the Josephson relation, which connects changes in the chemical potential to the time derivative of the phase, actually allows one to interpret the *compressibility*, κ , as a helicity modulus in the imaginary time direction. Thus κ enters a free energy correction $\Delta f_\tau = (\kappa/2\beta V) \int_0^\beta d\tau \int d^d x (\partial_\tau \phi)^2$. The same logic then suggests that $\partial_\tau \phi$ should scale as $\xi_\tau^{-1} \sim |\delta|^{z\nu}$, leading to $\kappa \sim |\delta|^{v_\tau}$ with $v_\tau = 2 - \alpha - 2z\nu = (d - z)\nu$. Since both the superfluid and the Bose glass phases have finite compressibility, one expects on physical grounds that κ should be finite and nonzero at $\delta = 0$ as well. This immediately leads to the prediction $z = d$ [2, 3]. However, a hidden assumption in these arguments is that Δf_x and Δf_τ can be included in the singular part of the free energy, f_s . We shall show that this is correct for Δf_x but not for Δf_τ . In fact the main contribution to Δf_τ comes from the nonsingular analytic part of the free energy, f_a , so that κ is dominated by its analytic part which is trivially finite at the transition, and f_s yields only small corrections to this that vanish at the critical point.

Rigorous definitions of the helicity moduli involve comparing free energies with twisted and untwisted boundary conditions. Previously such boundary condition dependence, Δf , has always been included in f_s : normally the strong dependence required for a finite helicity modulus requires long or quasi-long range order that is present only in the superfluid phase. The fact that $\Upsilon \equiv 0$ in the disordered phase then guarantees that it can arise only from singular terms in the free energy. In the present problem, however, κ is nonzero in both phases:

the gapless excitation spectrum of the Bose glass phase leads to power law order in imaginary time (though not in space) [3]. The free energy of both phases therefore have strong temporal boundary condition dependence, and an analytic contribution is actually very natural.

From our analysis there emerges the following general criterion for when Δf can be dominated by f_a . If the twisted boundary condition breaks a fundamental symmetry of the model, then it can be expected to generate relevant (in the renormalization group sense) terms in the Lagrangian. This leads to a new relevant, diverging scaling variable in the singular part of the free energy that generally will dominate all other contributions to Δf . In this case the usual Josephson scaling relation will hold. On the other hand, if no additional symmetry is broken, no new relevant scaling variable results and the twist will lead only to small shifts in the parameters already present in the untwisted Lagrangian. The helicity modulus, which involves derivatives with respect to these shifts, will then be dominated by f_a . In the present problem the twist couples to particle-hole symmetry [9] which is always broken at the Bose glass to superfluid transition.

For convenience we consider a continuum $|\psi|^4$ -model functional integral representation of the partition function, $Z = \int D\psi \exp(-\mathcal{L}_B)$, with Lagrangian [3, 9]

$$\begin{aligned} \mathcal{L}_B = & \int_0^\beta d\tau \int d^d x \left\{ -J\psi^*(\mathbf{x}, \tau) \nabla^2 \psi(\mathbf{x}, \tau) \right. \\ & - K\psi^*(\mathbf{x}, \tau) [\partial_\tau - \mu(\mathbf{x})]^2 \psi(\mathbf{x}, \tau) \\ & \left. + r(\mathbf{x})|\psi(\mathbf{x}, \tau)|^2 + u|\psi(\mathbf{x}, \tau)|^4 \right\}, \end{aligned} \quad (1)$$

where $J \approx \hbar^2/2m > 0$ is the boson hopping amplitude which favors spatial ferromagnetic order in the phase of the field ψ ; $\mu(\mathbf{x})$ is the (static) random external potential; and $K \approx 1/2u_0 > 0$, where u_0 is the soft core repulsion between nearby bosons, favors ferromagnetic order along the imaginary time dimension as well. This model is a continuum approximation to the Josephson junction array Lagrangian in which $\psi(\mathbf{x}, \tau) \rightarrow \exp[i\phi_i(\tau)]$ where $\phi_i(\tau)$ is the Josephson phase at site i , and the $\psi^* \nabla^2 \psi$ term is the continuum limit of the Josephson coupling, $J_{ij} \cos[\phi_i(\tau) - \phi_j(\tau)]$, between two nearby sites i and j . The $r|\psi|^2 + u|\psi|^4$ terms then represent the usual soft spin approximation to the the constraint $|\psi| = 1$. We allow $r(\mathbf{x})$ to be random as well, representing disorder in the hopping amplitudes, J_{ij} . We could also allow J and u to be random, but this is less convenient and does not produce anything new. We write $\mu(\mathbf{x}) = \mu - \epsilon(\mathbf{x})$ and $r(\mathbf{x}) = r_0 + w(\mathbf{x})$ with disorder average $[\epsilon(\mathbf{x})]_{av} = [w(\mathbf{x})]_{av} = 0$. For a given value of μ , the transition to superfluidity occurs with increasing J at a point $J_c(\mu)$, and we shall take $\delta = J - J_c(\mu)$.

Other representations of the problem serve equally well. The arguments we shall present are very general

and do not depend on the precise form of $\mathcal{L}_B$. As seen below, the only common feature required is that time derivatives and chemical potential always appear in the combination $(\partial_\tau - \mu)$. In the coherent state functional integral formulation, for example, the Lagrangian has only a linear term, $\psi^*(\partial_\tau - \mu)\psi$. The two formulations match up to irrelevant terms (in the renormalization group sense) so long as $\mu(\mathbf{x})$ does not vanish identically [3, 9]. When $\mu(\mathbf{x}) \equiv 0$ the model (1) reduces to the well known random rod problem, corresponding to a classical $(d+1)$ -dimensional XY -model with columnar disorder [9, 10, 11]. This model maintains an exact particle-hole symmetry, and the Josephson scaling relation for κ is therefore valid, but with $z < d$: see below.

Twisted θ -boundary conditions [12] are defined by the condition

$$\psi(x + L_\alpha \hat{\mathbf{e}}_\alpha) = e^{i\theta_\alpha} \psi(x), \quad |\theta_\alpha| \leq \pi, \quad (2)$$

where $x \equiv (\tau, \mathbf{x})$, $\hat{\mathbf{e}}_\alpha$, $\alpha = 0, 1, \dots, d$, are the space-time unit vectors, and L_α is the dimension of the system along $\hat{\mathbf{e}}_\alpha$ (with $L_0 = \beta$). Thus $\bar{\psi}(\mathbf{x}) = e^{-i\mathbf{k}_0 \cdot \mathbf{x} - i\omega_0 \tau} \psi(\mathbf{x})$, where $\omega_0 = \theta_0/\beta$ and $\mathbf{k}_0 = (\theta_1/L_1, \dots, \theta_d/L_d)$, obeys periodic boundary conditions. We therefore expect $\bar{\psi}_0 = \langle \bar{\psi}(\mathbf{x}) \rangle$ to be uniform and hence the order parameter $\langle \psi(\mathbf{x}) \rangle = e^{i\mathbf{k}_0 \cdot \mathbf{x} + i\omega_0 \tau} \bar{\psi}_0$ to have a uniform phase twist. We now rewrite the Lagrangian in terms of $\bar{\psi}$, obtaining

$$\mathcal{L}_B^{\omega_0 \mathbf{k}_0}[\bar{\psi}; \mu, r_0] = \mathcal{L}_B[\bar{\psi}; \mu - i\omega_0; r_0 + J\mathbf{k}_0^2] + \mathbf{k}_0 \cdot \mathbf{P}[\bar{\psi}]. \quad (3)$$

where $\mathbf{P}[\bar{\psi}] = -iJ \int d^d x \int_0^\beta d\tau [\bar{\psi}^* \nabla \bar{\psi} - \bar{\psi} \nabla \bar{\psi}^*]$ is the total momentum. The simple shift, $\mu \rightarrow \mu - i\omega_0$, is guaranteed by the combination $(\partial_\tau - \mu)$ for any model of dirty bosons.

The free energy is defined by $f^{\mathbf{k}_0 \omega_0} = (\beta V)^{-1} [\ln(Z)]_{av}$, where the superscripts label the boundary condition. Using (3), straightforward expansion of the free energy in powers of $\mathbf{k}_0$ at $\omega_0 = 0$,

$$\Delta f_x \equiv f^{\mathbf{k}_0} - f = \frac{1}{2} \Upsilon \mathbf{k}_0^2 + O(\mathbf{k}_0^4). \quad (4)$$

produces the usual momentum-momentum type correlation function expression for Υ . On the other hand, if $\mathbf{k}_0 = 0$ we have the obvious result $f^{\omega_0}(\mu) = f(\mu - i\omega_0)$. Analyticity of f within a given thermodynamic phase immediately implies the expansion

$$\Delta f_\tau \equiv f^{\omega_0} - f = i\omega_0 \rho + \frac{1}{2} \kappa \omega_0^2 + O(\omega_0^3), \quad (5)$$

where $\rho = -\partial f / \partial \mu$ and $\kappa = \partial \rho / \partial \mu$. It is the comparison of equations (4) and (5) that motivates the identification of κ with the temporal helicity modulus. Note, however, that the density ρ actually yields the leading term linear in ω_0 . The reason this term appears in (5) but not in (4) is that $\mathcal{L}_B$ contains only quadratic spatial derivatives and is therefore even under space inversion, $\mathbf{x} \rightarrow -\mathbf{x}$. On the other hand $\mathbf{P}[\bar{\psi}]$ is odd under space inversion, implying

that $f^{\mathbf{k}_0\omega_0}$ is an even function of $\mathbf{k}_0$. Moreover we see that $\mathbf{k}_0$ breaks the inversion symmetry of $\mathcal{L}_B[\psi]$. In contrast, only if $\mu(\mathbf{x}) \equiv 0$ does $\mathcal{L}_B$ possess inversion symmetry in time, $\tau \rightarrow -\tau$, which we call particle-hole symmetry [13]. This symmetry is already broken by μ , and the additional twist, ω_0 , produces nothing new. These observations will be crucial to understanding the scaling of Υ and κ .

Let us consider first the Josephson scaling argument for the random rod problem, which then possesses both inversion and particle-hole symmetry. Classical intuition suggests in addition that $\kappa \equiv 0$ in the disordered phase [14]. As alluded to earlier, we then expect the leading boundary condition dependence to enter via singular scaling combinations $k_0\xi$ and $\omega_0\xi_\tau$. Formally, this expectation is implemented via a finite-size scaling ansatz for the free energy density [3, 12],

$$\Delta f^{\mathbf{k}_0\omega_0} \approx \beta^{-1} L^{-d} \Phi_0^{\mathbf{k}_0\omega_0} (A_0 \delta L^{1/\nu_0}, B_0 \delta \beta^{1/z_0\nu_0}), \quad (6)$$

where A_0 and B_0 are nonuniversal scale factors [15]. For convenience we take a hypercubical volume $V = L^d$, and the exponents are those of the classical random rod problem. The existence of a nonzero stiffness, i.e., a leading finite-size correction of order L^{-2} or β^{-2} , now requires that the scaling function $\Phi_0^{\mathbf{k}_0\omega_0}(x, y) \approx x^{d\nu_0} y^{z_0\nu_0} (Q_x x^{-2\nu_0} + Q_\tau y^{-2z_0\nu_0})$ for large $x, y > 0$, yielding

$$\begin{aligned} \Upsilon &\approx A_0^{(d-2)\nu_0} B_0^{z_0\nu_0} (Q_x/\theta_x^2) \delta^{\nu_0} \\ \kappa &\approx A_0^{d\nu_0} B_0^{-z_0\nu_0} (Q_0/\theta_0^2) \delta^{v_{\tau 0}}, \end{aligned} \quad (7)$$

where $\theta_x^2 = \theta_1^2 + \dots + \theta_d^2$, implying the Josephson scaling relations $\nu_0 = (d + z_0 - 2)\nu_0 = 2 - \alpha_0 - 2\nu_0$ and $v_{\tau 0} = (d - z_0)\nu_0 = 2 - \alpha_0 - 2z_0\nu_0$, and requiring in addition $Q_{x,0} \propto \theta_{x,0}^2$.

Consider now the Bose glass to superfluid transition. The density and compressibility of both phases are smooth nontrivial functions of μ , and a temporal twist now perturbs only slightly the particle-hole symmetry breaking term that is already in $\mathcal{L}_B$. On the other hand, spatial twists still produce singular corrections, and at $\omega_0 = 0$ we predict a finite-size scaling form [16]

$$\Delta f^{\mathbf{k}_0} = \beta^{-1} L^{-d} \Phi^{\mathbf{k}_0} (A \delta L^{1/\nu}, B \delta \beta^{1/z\nu}), \quad (8)$$

with $\Phi^{\mathbf{k}_0}(x, y) \approx R_x x^{(d-2)\nu} y^{z\nu}$ for large $x, y > 0$, yielding $\Upsilon \approx A^{(d-2)\nu} B^{z\nu} (R_x/\theta_x^2) \delta^v$, $v = (d + z - 2)\nu = 2 - \alpha - 2\nu$, and $R_x \propto \theta_x^2$ as before. All exponents now refer to the dirty boson critical point. Because we have, as yet, imposed no temporal twist, *there can be no* $O(\beta^{-1}, \beta^{-2})$ terms.

Now, if a finite ω_0 is included, the only changes in (8) are that $\mu \rightarrow \mu - i\omega_0$ everywhere, and boundary condition dependence of f_a must be included. Thus

$$\begin{aligned} \Delta f^{\mathbf{k}_0\omega_0} &= \beta^{-1} L^{-d} \Phi^{\mathbf{k}_0} (A \delta_\theta L^{1/\nu}, B \delta_\theta \beta^{1/z\nu}) \\ &\quad + f_a(J, r_0 + J\mathbf{k}_0^2, \mu - i\omega_0) - f_a(J, r_0, \mu), \end{aligned} \quad (9)$$

where $\delta_\theta = J - J_c(\mu - i\omega_0) \approx \delta + i\omega_0 J'_c(\mu)$. Most importantly, $\Phi^{\mathbf{k}_0}$ is the same function as that in (8), and therefore still yields no $O(\beta^{-1}, \beta^{-2})$ terms. This implies immediately that the scaling function $\Phi^{\mathbf{k}_0\omega_0}$ produces no *direct* contributions to ρ and κ , which must therefore arise (a) *indirectly* from the ω_0 dependence of δ_θ , and (b) directly from the analytic part of the free energy. The former couples derivatives with respect to μ , or equivalently ω_0 , to derivatives with respect to δ , producing the leading singular terms, $\kappa_s \sim |\delta|^{-\alpha}$ [17]. However, $\alpha = 2 - (d+z)\nu$ is very likely negative [3], so this gives a vanishing contribution at the critical point. Therefore the main contribution is analytic in origin. Taking $\mathbf{k}_0 = 0$, we may write $f_a(J, \mu) = -\rho_c(J)[\mu - \mu_c(J)] - \frac{1}{2}\kappa_c(J)[\mu - \mu_c(J)]^2 + \dots$, expanded for convenience about the transition line $\mu_c(J)$, and we immediately obtain from (9) a finite compressibility right through the transition, with the exponent z nevertheless *undetermined* [18, 19].

There is a second approach to the derivation of the Josephson scaling relation, through scaling of the two point correlation function in the superfluid hydrodynamic regime, which we now discuss. Long wavelength, low frequency fluctuations are known to be governed by the effective Gaussian Lagrangian [2, 3]

$$\mathcal{L}_{\text{HD}} = \frac{1}{2} \int d^d x \int_0^\beta d\tau [\Upsilon |\nabla \tilde{\phi}|^2 + \kappa (\partial_\tau \tilde{\phi})^2], \quad (10)$$

where $\tilde{\phi}(\mathbf{x}, \tau)$ is the coarse-grained phase, related to the coarse grained order parameter field via $\tilde{\psi}(\mathbf{x}, \tau) = \psi_0 \exp[i\tilde{\phi}(\mathbf{x}, \tau)]$, where the bulk order parameter, $\psi_0 \sim |\delta|^\beta$, near criticality. The small $\mathbf{k}, \omega$ form of the Fourier transform of the two point correlator, $G(\mathbf{x} - \mathbf{x}', \tau - \tau') = [\langle \psi^*(\mathbf{x}, \tau) \psi(\mathbf{x}', \tau') \rangle]_{av}$, is then

$$G(\mathbf{k}, \omega) \approx |\psi_0|^2 / [\Upsilon \mathbf{k}^2 + \kappa \omega^2]. \quad (11)$$

Normally it is assumed that G obeys the scaling form

$$G(\mathbf{k}, \omega) \approx C |\delta|^{-\gamma} g(Dk|\delta|^{-\nu}, E\omega|\delta|^{-z\nu}), \quad (12)$$

where γ is the susceptibility exponent and C, D, E are again nonuniversal scale factors. If one naively matches (11) and (12) one concludes that $g(x, y) \approx (g_1 x^2 + g_2 y^2)^{-1}$ for small x, y , and hence that

$$\begin{aligned} \Upsilon/|\psi_0|^2 &\approx g_1 C^{-1} D^2 |\delta|^{\gamma-2\nu} \\ \kappa/|\psi_0|^2 &\approx g_2 C^{-1} E^2 |\delta|^{\gamma-2z\nu}. \end{aligned} \quad (13)$$

Using the scaling relation $\alpha + 2\beta + \gamma = 2$ we obtain again the Josephson relations $v = 2 - \alpha - 2\nu$, $v_\tau = 2 - \alpha - 2z\nu$.

This argument, however, is just a disguised version of the free energy argument: $\mathcal{L}_{\text{HD}}$ is obtained by assuming that the energetics describing global phase twists also describes slowly varying local phase twists. Thus, locally we replace $\mathbf{k}_0$ by $\nabla \tilde{\phi}$ and ω_0 by $\partial_\tau \tilde{\phi}$, then integrate over all of space-time [20]. Since κ arises from the nonscaling

part of the free energy, it is unlikely that it can now arise from the scaling part of the two-point function. Rather, we must carefully reconsider the scaling ansatz for G . Our proposal (for which we have no detailed theoretical support at this stage), is that

$$|\psi_0|^2/G(\mathbf{k}, \omega) \approx H|\delta|^{2-\alpha}\Gamma(Dk|\delta|^{-\nu}, E\omega|\delta|^{-z\nu}) + \Gamma_a(\mathbf{k}, \omega), \quad (14)$$

where Γ_a is analytic. This self-energy scaling form is very similar to (9). Matching with (11), we again assume that Υ arises from the scaling part, κ from the analytic part. Thus $\Gamma(x, y) \approx \gamma_1 x^2 + \gamma_2 y^2$, for small x, y while $\Gamma_a(k, \omega) \approx \kappa \omega^2$ for small k, ω . The x^2 term yields $\Upsilon \sim |\delta|^\nu$. If $z < d$ the y^2 term is subdominant to the analytic term, which yields κ finite, as required, leaving z undetermined [21, 22]. If we consider only static correlations, $\omega = 0$, standard scaling is recovered without any unusual analytic corrections.

To summarize, the original scaling ansatz for the Bose glass-superfluid critical point, in which temporal twists produce a relevant symmetry breaking perturbation to the Lagrangian, and therefore scale in the combination $\omega_0 \xi_\tau$, is not supported theoretically, and may explain violations of $z = d$ in $d = 2$ [8].

Acknowledgments: We thank D. S. Fisher for fruitful discussions.

-
- [1] M. Ma, B. I. Halperin and P. A. Lee, Phys. Rev. B **34**, 3136 (1986).
[2] D. S. Fisher and M. P. A. Fisher, Phys. Rev. Lett. **61**, 1847 (1988).
[3] M. P. A. Fisher, P. B. Weichman, G. Grinstein and D. S. Fisher, Phys. Rev. B **40**, 546 (1989).
[4] P. B. Weichman and K. Kim, Phys. Rev. B **40**, 813, (1989).
[5] I. F. Herbut, Phys. Rev. B **57**, 1303 (1998).
[6] See, e.g., R. T. Scalettar, G. G. Batrouni and G. T. Zimanyi, Phys. Rev. Lett. **66**, 3144 (1991); S. Zhang, N. Kawashima, J. Carlson and J. E. Gubernatis, Phys. Rev. Lett. **74**, 1500 (1995); F. Alet and E. S. Sørensen, Phys. Rev. E **67**, 015701(R) (2003); N. Prokof'ev and B. Svistunov, Phys. Rev. Lett. **92**, 015703 (2004).
[7] M. Makivic, N. Trivedi and S. Ullah, Phys. Rev. Lett. **71**, 2307 (1993); P. B. Weichman, Phys. Rev. Lett. **74**, 1038 (1995); N. Trivedi and M. Makivic, Phys. Rev. Lett. **74**, 1039, (1995).
[8] A. Priyadarshree, S. Chandrasekharan, J.-W. Lee, and H. U. Baranger, Phys. Rev. Lett. **97**, 115703 (2006).
[9] R. Mukhopadhyay and P. B. Weichman, Phys. Rev. Lett. **76**, 2977 (1996).
[10] S. N. Dorogovtsev, Phys. Lett. **76A**, 169 (1980); D. Boyanovsky and J. L. Cardy, Phys. Rev. B **26**, 154 (1982).
[11] P. B. Weichman and R. Mukhopadhyay (unpublished).
[12] M. E. Fisher, M. N. Barber and D. Jasnow, Phys. Rev. A **8**, 1111 (1973). Amazingly, the last section of this paper, containing a discussion of how finite size scaling might lead to violations of the Josephson relation, proposes a mechanism very similar to ours: anomalously strong boundary condition dependence of t_θ , the distance from the critical point, on the twist wavevector, k_0 . If $t_\theta - t \sim k_0^2$ they find $\nu = 1 - \alpha$. Here we find an even more anomalous linear dependence of $\delta_\theta - \delta$ on ω_0 , leading to $\nu_\tau = -\alpha$. We are unaware of any classical model where the original proposed violation occurs.
[13] This must be contrasted with time reversal symmetry, which involves a simultaneous replacement $\psi \leftrightarrow \psi^*$, which $\mathcal{L}_B$ always satisfies.
[14] This can be demonstrated more explicitly: P. B. Weichman and R. Mukhopadhyay (unpublished).
[15] No overall multiplicative scale factor is required due to quantum hyperuniversality: see K. Kim and P. B. Weichman, Phys. Rev. B **43**, 13583 (1991).
[16] In Ref. [9] it was argued that so long as $\epsilon(\mathbf{x}) \neq 0$, μ itself was irrelevant, i.e., a statistical particle-hole symmetry is restored at the critical point. This would imply a correction to scaling combination $\mu|\delta|^{\lambda_\mu}$, $\lambda_\mu > 0$ which we have ignored in (8). This means that μ drops out completely sufficiently close to the critical point, ruling out again any scaling of chemical potential changes with ξ_τ .
[17] Within the ϵ, ϵ_τ -expansion [10], where ϵ_τ is the dimension of time and $\epsilon = 4 - d - \epsilon_\tau$, it was found [9] that the Bose glass phase is compressible *only* in the physical dimension, $\epsilon_\tau = 1$, with $\kappa \equiv 0$ for any $\epsilon_\tau < 1$. This precludes any analytic contribution to κ . We expect then that $\kappa \sim |\delta|^{-\alpha(\epsilon, \epsilon_\tau)} \rightarrow 0$ from the superfluid side.
[18] The fact that the leading correction is *linear* in ω_0 leads even more directly to problems with the previous scaling ansatz [2, 3]: if a finite $\kappa = \partial\rho/\partial\mu$ is to be associated with the singular part of the free energy, it must come from the singular part of the density which, for consistency, must scale as $\xi^{-d} \sim |\delta|^{d\nu}$. This vanishes rapidly at the critical point, contradicting the original assumption that the transition takes place at finite boson density.
[19] This can be confirmed explicitly in $d = 1$ [14]. Except for rapidly decaying irrelevant terms, μ can be removed entirely from the dual surface roughening model Lagrangian via a shift, appearing, consistent with our general arguments, only as an additive analytic term in the free energy. The result $z = 1$ for this model is therefore unconnected to the appearance of a finite κ .
[20] The linear term, $i\rho(\partial_\tau\phi)$ integrates to an integer multiple of $2\pi i\rho$ and disappears from $\exp(-\mathcal{L}_{\text{HD}})$ since the total boson number ρV is an integer.
[21] There are dynamical scaling breakdown issues at the finite T lambda transition (described by the classical Model F equations) as well, where the finite κ argument now yields $z = d/2$. However, violations are possible where there exist two different dynamical exponents z_1, z_2 satisfying $z_1 + z_2 = d$. Only the mean $(z_1 + z_2)/2 = d/2$ enters the corresponding hydrodynamic correlation function [V. Dohm, Phys. Rev. B **44** 2697 (1991)].
[22] In a perhaps related effect, the rigorous finite-size correlation length exponent inequality $\nu_{\text{FS}} \geq 2/d$ for classical random systems [J. T. Chayes, L. Chayes, D. S. Fisher, and T. Spencer, Phys. Rev. Lett. **57**, 2999 (1986)] does not preclude a second “intrinsic” exponent ν with $\nu < 2/d$ [F. Pázmándi, R. Scalettar and G. T. Zimányi, Phys. Rev. Lett. **79**, 5130 (1997)].